

# Higher Dimensional Transition System and Labelled Symmetric Precubical Set

*(talk in French, slides in English)*

Philippe Gaucher



<http://www.pps.jussieu.fr/~gaucher>

Preuves, Programmes et Systèmes, CNRS UMR 7126 et Paris 7

# Organization of the talk

1. Weak higher dimensional transition system (HDTS)
2. Labelled symmetric precubical set
3. Realizing a labelled symmetric precubical set as a weak HDTS
4. Categorical equivalence
5. Homotopy theory of weak HDTS

# Concurrent execution of two actions



- $\{\alpha, \beta, \gamma, \delta\}$  set of states and of 0-cubes
- $\{(\alpha, a, \beta), (\beta, b, \gamma), (\alpha, b, \delta), (\delta, a, \gamma), (\alpha, a, b, \beta), (\alpha, b, a, \beta)\}$  set of transitions

# Weak HDTS


- $\Sigma$  a nonempty set of labels
- A weak HDTS  $X = (S, \mu : L \rightarrow \Sigma, T = \bigcup_{n \geq 1} T_n)$  with set of states  $S$ , set of actions  $L$ , labelling map  $\mu$ , set of  $n$ -transitions  $T_n \subset S \times L^n \times S$  with  $n \geq 1$ 
  - **Multiset axiom**  $(\alpha, u_1, \dots, u_n, \beta) \in T$  implies  $(\alpha, u_{\sigma(1)}, \dots, u_{\sigma(n)}, \beta) \in T$  for every permutation  $\sigma$
  - **Coherence axiom** if  $(\alpha, u_1, \dots, u_n, \beta)$ ,  $(\alpha, u_1, \dots, u_p, \nu_1)$ ,  $(\nu_1, u_{p+1}, \dots, u_n, \beta)$ ,  $(\alpha, u_1, \dots, u_{p+q}, \nu_2)$  and  $(\nu_2, u_{p+q+1}, \dots, u_n, \beta)$  belong to  $T$ , then  $(\nu_1, u_{p+1}, \dots, u_{p+q}, \nu_2) \in T$
- Note: the Coherence axiom is automatically satisfied in a cube

# Category WHDTS

- $X = (S, \mu : L \rightarrow \Sigma, T = \bigcup_{n \geq 1} T_n)$  and  
 $X' = (S', \mu' : L' \rightarrow \Sigma, T' = \bigcup_{n \geq 1} T'_n)$
- A map of weak HDTS  $f : X \rightarrow X'$  consists of
  - a set map  $f_0 : S \rightarrow S'$
  - a map  $\tilde{f} \in (\mathbf{Set} \downarrow \Sigma)(\mu, \mu')$such that if  $(\alpha, u_1, \dots, u_n, \beta)$  is a transition, then  $(f_0(\alpha), \tilde{f}(u_1), \dots, \tilde{f}(u_n), f_0(\beta))$  is a transition
- The forgetful functor  $\omega : \mathbf{WHDTS} \rightarrow \mathbf{Set}^{\{s\} \cup \Sigma} :$   
 $\omega(X) = (S, (\mu^{-1}(x))_{x \in \Sigma})$  is **concrete topological**
- WHDTS is **locally finitely presentable**

# Two additional axioms

- **First Cattani-Sassone Axiom CSA1** If  $(\alpha, u, \beta)$  and  $(\alpha, u', \beta)$  are two transitions such that  $\mu(u) = \mu(u')$ , then



- CSA1 always satisfied for concrete examples
- **Unique Intermediate State Axiom CSA2** for  $n \geq 2$  and  $1 \leq p < n$ , for every transition  $(\alpha, u_1, \dots, u_n, \beta)$ , there exists a unique state  $\nu$  such that  $(\alpha, u_1, \dots, u_p, \nu)$  and  $(\nu, u_{p+1}, \dots, u_n, \beta)$  are transitions
  - CSA2 plays the role of the face maps in a precubical set

# The category HDTS

- HDTS : weak HDTS satisfying CSA1 and CSA2
  - HDTS **locally finitely presentable**, but **not topological**
  - HDTS full **reflective** subcategory of WHDTS
- The **pure  $n$ -transition**  $C_n[a_1, \dots, a_n]^{ext}$ 
  - Set of states  $S = \{I, F\}$
  - Set of actions  $L = \{(a_1, 1), \dots, (a_n, n)\}$
  - Labelling map  $\mu(a_i, i) = a_i$
  - $T = \{(I, (a_{\sigma(1)}, \sigma(1)), \dots, (a_{\sigma(n)}, \sigma(n)), F)\}$
- The  **$n$ -cube**  $C_n[a_1, \dots, a_n]$  is the reflection of  $C_n[a_1, \dots, a_n]^{ext}$ :  $2^n$  states since  $\{(a_1, 1), \dots, (a_n, n)\}$  contains  $n$  elements

# Cattani-Sassone HDTS

- A Cattani-Sassone HDTS

$X = (S, \mu : L \rightarrow \Sigma, T = \bigcup_{n \geq 1} T_n)$  with set of states  $S$ , set of actions  $L$ , labelling map  $\mu$ , set of  $n$ -transitions  $T_n \subset S \times L^n \times S$  with  $n \geq 1$

- Multiset axiom, CSA1, CSA2

- **Third Cattani-Sassone axiom CSA3** if the nine

tuples  $(\alpha, u_1, \dots, u_n, \beta)$ ,  $(\alpha, u_1, \dots, u_p, \nu_1)$ ,

$(\nu_1, u_{p+1}, \dots, u_n, \beta)$ ,  $(\nu_1, u_{p+1}, \dots, u_{p+q}, \nu_2)$ ,

$(\nu_2, u_{p+q+1}, \dots, u_n, \beta)$ ,  $(\alpha, u_1, \dots, u_{p+q}, \nu'_2)$ ,

$(\nu'_2, u_{p+q+1}, \dots, u_n, \beta)$ ,  $(\alpha, u_1, \dots, u_p, \nu'_1)$  and

$(\nu'_1, u_{p+1}, \dots, u_{p+q}, \nu'_2)$  are transitions, then  $\nu_1 = \nu'_1$

and  $\nu_2 = \nu'_2$

- The HDTS are the Cattani-Sassone HDTS

# Symmetric precubical set

## ● Symmetric precubical set:

- Family  $(K_n)_{n \geq 0}$  of sets ( $x \in K_n$  is called a  $n$ -cube), of **face maps**  $\partial_i^\alpha : K_n \rightarrow K_{n-1}$  satisfying the cubical relations and of **symmetry maps**  $s_i : K_n \rightarrow K_n$  with  $1 \leq i \leq n - 1$  satisfying the Moore relations and  $\partial_j^\alpha s_i = s_{i-1} \partial_j^\alpha$  for  $j < i$ ,  $\partial_j^\alpha s_i = \partial_{i+1}^\alpha$  for  $j = i$ ,  $\partial_j^\alpha s_i = \partial_i^\alpha$  for  $j = i + 1$  and  $\partial_j^\alpha s_i = s_i \partial_j^\alpha$  for  $j > i + 1$

## ● The symmetric precubical set of labels:

- $(!^S \Sigma)_0 = \{()\}$
- $(!^S \Sigma)_n = \Sigma^n$  for  $n \geq 1$
- $\partial_i^\alpha(a_1, \dots, a_n) = (a_1, \dots, \hat{a}_i, \dots, a_n)$
- $s_i(a_1, \dots, a_n) = (a_1, \dots, a_{i-1}, a_{i+1}, a_i, a_{i+2}, a_n)$

# Labelled symmetric precubical set

- **Labelled symmetric precubical set** : map  $K \rightarrow !^S \Sigma$ 
  - The category  $\square_S^{op} \text{Set} \downarrow !^S \Sigma$  is **locally finitely presentable**
- $[n] = \{0, 1\}^n$ , the  **$n$ -cube**  $\square_S[n]$  defined by:  $f \in \square_S[n]_p$  set map from  $[p]$  to  $[n]$  which is a composite of
  - $\delta_i^\alpha(\epsilon_1, \dots, \epsilon_r) = (\epsilon_1, \dots, [\alpha]_i, \dots, \epsilon_r)$
  - $\sigma_i(\epsilon_1, \dots, \epsilon_r) = (\epsilon_1, \dots, \epsilon_{i+1}, \epsilon_i, \dots, \epsilon_r)$
- The **labelled  $n$ -cube**  $\square_S[a_1, \dots, a_n]$  with  $a_1, \dots, a_n \in \Sigma$  is the map  $\square_S[n] \rightarrow !^S \Sigma$  which takes  $\text{Id}_{[n]}$  to  $(a_1, \dots, a_n)$
- The **boundary**  $\partial \square_S[a_1, \dots, a_n]$  of  $\square_S[a_1, \dots, a_n]$  is the labelled  $n$ -cube with the  $n!$   $n$ -cubes removed

# From precubical set to WHDTS (I)

- There exists an **isomorphism of categories**

$$\mathbb{T} : \left\{ \text{Cubes of } \square_S^{op} \text{Set} \downarrow !^S \Sigma \right\} \cong \left\{ \text{Cubes of WHDTS} \right\}$$

- $\mathbb{T}$  extended to a functor  $\mathbb{T} : \square_S^{op} \text{Set} \downarrow !^S \Sigma \rightarrow \mathbf{WHDTS}$  by

$$\mathbb{T}(K) := \lim_{\square_S[a_1, \dots, a_n] \rightarrow K} C_n[a_1, \dots, a_n]$$

- $\mathbb{T} : \square_S^{op} \text{Set} \downarrow !^S \Sigma \rightarrow \mathbf{WHDTS}$  is a **left adjoint**
- $\mathbb{T}$  is not faithful, not full and not essentially surjective

# From precubical set to WHDTS (II)

- $\mathbb{T}(\square_S[a_1, \dots, a_m]) = C_m[a_1, \dots, a_m]$
- For  $f : \square_S[a_1, \dots, a_m] \rightarrow \square_S[b_1, \dots, b_n]$ ,  
 $\mathbb{T}(f) : C_m[a_1, \dots, a_m] \rightarrow C_n[b_1, \dots, b_n]$  is the pair  
 $((\mathbb{T}(f))_0, \widetilde{\mathbb{T}(f)})$ 
  - $(\mathbb{T}(f))_0 = f_0 : [m] \rightarrow [n]$
  - $\widetilde{\mathbb{T}(f)}(a_i, i) = (a_i, \overline{f}^{-1}(i)) = (b_{\overline{f}^{-1}(i)}, \overline{f}^{-1}(i))$

with

- $f_0(\epsilon_1, \dots, \epsilon_m) = (\epsilon_{\widehat{f}(1)}, \dots, \epsilon_{\widehat{f}(n)})$
- $\widehat{f} : \{1, \dots, n\} \rightarrow \{1, \dots, m\} \cup \{-\infty, +\infty\}$  with  
 $\epsilon_{-\infty} = 0$  and  $\epsilon_{+\infty} = 1$
- $\overline{f} : \widehat{f}^{-1}(\{1, \dots, m\}) \rightarrow \{1, \dots, m\}$  **bijective**

# About faithfulness

- For  $n \geq 2$ , the two maps

$$\square_S[a_1, \dots, a_n] \rightrightarrows \square_S[a_1, \dots, a_n] \sqcup_{\partial \square_S[a_1, \dots, a_n]} \square_S[a_1, \dots, a_n]$$

have same image by  $\mathbb{T}$

$$\text{Id}_{C_n[a_1, \dots, a_n]} : C_n[a_1, \dots, a_n] \rightarrow C_n[a_1, \dots, a_n]$$

- The map  $(\square_S^{op} \text{Set} \downarrow !^S \Sigma)(K, L) \rightarrow \mathbf{WHDTs}(\mathbb{T}(K), \mathbb{T}(L))$  is **one-to-one** when there exists at most one lift  $k$

$$\begin{array}{ccc} \partial \square_S[a_1, \dots, a_n] & \xrightarrow{\quad} & L \\ \downarrow & \nearrow k & \\ \square_S[a_1, \dots, a_n] & & \end{array}$$

for  $n \geq 2$  and  $a_1, \dots, a_n \in \Sigma$  **HDA paradigm**

# About the HDA paradigm

- A labelled symmetric precubical set  $L$  satisfies the **HDA paradigm** if and only if  $L$  is **orthogonal** to the maps


$$\square_S[a_1, \dots, a_n] \sqcup_{\partial \square_S[a_1, \dots, a_n]} \square_S[a_1, \dots, a_n] \rightarrow \square_S[a_1, \dots, a_n]$$

for  $n \geq 2$  and  $a_1, \dots, a_n \in \Sigma$

- The full subcategory  $\mathbf{HDA}^\Sigma$  of  $L$  satisfying the HDA paradigm is **reflective locally finitely presentable**
- The restriction functor  $\mathbb{T} : \mathbf{HDA}^\Sigma \rightarrow \mathbf{WHDTs}$  is **faithful**
- It is not full, nor essentially surjective...

# About fullness (I)

- Let  $C_2[u, v]$  be the HDTS with set of states  $\{\alpha, \beta, \nu_0, \nu_2\}$
- Let  $X$  be the image by  $\mathbb{T}$  of



- The inclusion  $C_2[u, v] \subset X$  does not come from a map of labelled symmetric precubical sets since there are no squares with the vertices  $\alpha, \beta, \nu_0, \nu_2$

# About fullness (II)

- Let  $K$  and  $L$  be two labelled symmetric precubical sets such that  $L$  satisfies the **HDA paradigm** and such that  $\mathbb{T}(L)$  satisfies the **Unique intermediate state axiom**. Then the set map

$$(\square_S^{op} \mathbf{Set} \downarrow !^S \Sigma)(K, L) \xrightarrow{f \mapsto \mathbb{T}(f)} \mathbf{WHDTS}(\mathbb{T}(K), \mathbb{T}(L))$$

is **bijjective**

# About essential surjectivity

- $\underline{x} = (\emptyset, \{x\} \subset \Sigma, \emptyset)$  with  $x \in \Sigma$  does not belong to the image of  $\mathbb{T}$ 
  - $\mathbf{HDTS}_{opt} := \{\underline{x} \subset C_1[x], x \in \Sigma\} - \text{inj} \subset \mathbf{HDTS}$  full subcategory of **optimized HDTS**: **locally finitely presentable and coreflective**
  - The right adjoint  $\text{Opt} : \mathbf{HDTS} \rightarrow \mathbf{HDTS}_{opt}$  removes all unused actions:  $\text{Opt}(\underline{x}) = \emptyset$
- The adjunction  $\subset : \mathbf{HDTS}_{opt} \rightleftarrows \mathbf{HDTS} : \text{Opt}$  gives rise to a category equivalence  $\mathbf{HDTS}_{opt} \simeq \mathbf{HDTS}[\text{Opt}^{-1}]$
- In  $\mathbf{HDTS}[\text{Opt}^{-1}]$ , two higher dimensional transition systems are isomorphic if they are isomorphic modulo their unused actions

# The category equivalence

- a category equivalence

$$\mathbf{HDA}_{hdt_s}^{\Sigma} := \mathbb{T}^{-1}(\mathbf{HDTS}) \cap \mathbf{HDA}^{\Sigma} \simeq \mathbf{HDTS}_{opt} [\mathcal{O}^{-1}]$$

where  $\mathcal{O}$  is the class of maps  $X \rightarrow Y$  inducing a bijection between the states, a surjection between the actions and such that  $Y$  is final:  $\varinjlim (C_1[x] \leftarrow \underline{x} \rightarrow C_1[x])$  is not of the form  $\mathbb{T}(K)$


- $\mathbf{HDA}_{hdt_s}^{\Sigma}$  is **reflective locally finitely presentable** in  $\mathbf{HDA}^{\Sigma}$ , and in  $\square_S^{op} \mathbf{Set} \downarrow !^S \Sigma$
- Note: a HDTS  $\mathbb{T}(K)$  satisfies CSA1 if and only if  $K$  is orthogonal to the maps  $\square_S[a] \sqcup_{\partial \square_S[a]} \square_S[a] \rightarrow \square_S[a]$  for  $a \in \Sigma$

# Homotopy theory of weak HDTs

- A **cofibration** of weak HDTs is a map inducing an injection on the set of actions
- There exists a **left-determined model category structure** on WHDTs with respect to the class of cofibrations
  - i.e. the smallest localizer  $\mathcal{W}$  with respect to the cofibrations is the class of weak equivalences of a model category structure
  - A left proper combinatorial model category
- Consider the **Bousfield-localization** with respect to the inclusions
  - $C_n[a_1, \dots, a_n]^{ext} \subset C_n[a_1, \dots, a_n]$
  - $\underline{x} \subset C_1[x]$

# Homotopy and bisimulation

- $\mathcal{P}$  the class of **paths**, i.e. of weak HDTS with a **unique initial state**  $I$  and a **unique final state**  $F$



- Two weak HDTS  $X$  and  $X'$  are  **$\mathcal{P}$ -bisimilar** if there exists a zig-zag of maps  $X \leftarrow Z \rightarrow X'$  such that  $Z \rightarrow X$  and  $Z \rightarrow X'$  satisfy the RLP with respect to the inclusion  $\{I\} \subset P$  for every  $P \in \mathcal{P}$
- Two weakly equivalent HDTS are  $\mathcal{P}$ -bisimilar