

Bar recursion in classical realisability : dependent choice and continuum hypothesis

Jean-Louis Krivine

May 16, 2016

Introduction

This paper is about the bar recursion operator [9], in the context of classical realizability [6, 7]. It is a sequel to the three papers [1, 2, 10]. We use the definitions and notations of the theory of classical realizability as expounded in [5, 6, 7].¹

In [1], S. Berardi, M. Bezem and T. Coquand have shown that a form of the bar recursion operator can be used, in a proof-program correspondence, to interpret *the axiom of dependent choice* in proofs of Π_2^0 -formulas of arithmetic. Their work was enhanced and adapted to the theory of domains by U. Berger and P. Oliva in [2]. In [10], T. Streicher has shown, by using the bar recursion operator of [2], that the models of ZF, associated with realizability algebras [5, 7] obtained from usual models of λ -calculus (Scott domains, coherent spaces, ...), satisfy the axiom of dependent choice.

We give here a proof of this result, but for a realizability algebra which is built following the presentation of [1], which we call the BBC-algebra.

In section 1, we define and study this algebra ; we define also the bar recursion operator, which is a closed λ -term.

In sections 2 and 3, which are very similar, we show that this operator realizes the axiom of countable choice (CC), then the axiom of dependent choice (DC). The proof is a little simpler for CC.

In section 4, we deduce from this result, using results of [8] that, in the model of ZF associated with this realizability algebra, *every real (more generally, every sequence of ordinals) is constructible*.

The formulas “ \mathbb{R} is well ordered” and “*Continuum Hypothesis*” are therefore realized in these models by a closed λ_c -term (i.e. a λ -term containing the control instruction cc of Felleisen-Griffin).

We show also that *every true formula of analysis is realized by a closed λ_c -term*.

Using these new results, we show how to obtain a program (closed λ_c -term) from any proof of a Π_2^0 arithmetical formula in the theory ZF + “Dependent choice” + “Every real is constructible” (and therefore “Well ordering of \mathbb{R} ” and “Continuum Hypothesis”).

¹Articles [5, 6, 7, 8] are available at www.pps.univ-paris-diderot.fr/~krivine/

1 The BBC realizability algebra

The definition and general properties of a *realizability algebra* are given at the beginning of [5]. It consists of a set of *terms* Λ , a set of *stacks* Π and a set of *processes* $\Lambda \star \Pi$. Every closed λ -term is interpreted as a term.

In this paper, we consider a particular realizability algebra $\mathcal{B} = (\Lambda, \Pi, \perp)$, which we call the *BBC algebra* because it is a reformulation of the programming language of [1]. It is defined as follows :

- The set of *processes* $\Lambda \star \Pi$ is $\Lambda \times \Pi$.
- The set of *terms* Λ is the smallest set which contains the following *constants of term* :
 B, C, I, K, W (*Curry's combinators*), cc (*Felleisen-Griffin instruction*), A (*abort instruction*),
 p, q_0, \dots, q_N (*variables*) where N is a fixed integer ;

and is such that :

if $\xi, \eta \in \Lambda$ then $(\xi)\eta \in \Lambda$ (*application*) ;
with each sequence $\xi_i (i \in \mathbb{N})$ of closed elements of Λ (i.e. which contain no variable p, q_0, \dots, q_N) is associated, in a one-to-one (and well founded) way, a constant of term denoted by $\bigwedge_i \xi_i$.

Therefore, each term $\xi \in \Lambda$ is a finite sequence of constants of term and parentheses. Λ is defined by an induction of length \aleph_1 and is of cardinality 2^{\aleph_0} .

Notations. The application $(\dots((\xi_1)\xi_2)\dots)\xi_n$ is often written $(\xi_1)\xi_2 \dots \xi_n$ or even $\xi_1 \xi_2 \dots \xi_n$. The finite sequence q_0, \dots, q_N will be often written \vec{q} .

- The set of *stacks* Π is defined as follows : a stack π is a finite sequence $t_0 \bullet \dots \bullet t_{n-1} \bullet \pi_0$ with $t_0, \dots, t_{n-1} \in \Lambda$; it is terminated by the symbol π_0 which represents the *empty stack*.

For each stack π , the *continuation* k_π is a term which is defined by recurrence :

$k_{\pi_0} = A$; $k_{t \bullet \pi} = \ell_t k_\pi$, with $\ell_t = ((C)(B)CB)t$ or $\lambda k \lambda x(k)(x)t$.

Thus, if the stack π is $t_0 \bullet \dots \bullet t_{n-1} \bullet \pi_0$, we have $k_\pi = (\ell_{t_0}) \dots (\ell_{t_{n-1}})A$ or $\lambda x(A)(x)t_0 \dots t_{n-1}$.

The *integer* \underline{n} is defined as follows :

$\underline{0} = (K)I$ or $\lambda x \lambda y y$; $\underline{n+1} = (\sigma)\underline{n}$ with $\sigma = (BW)(C)(B)BB$ or $\lambda n \lambda f \lambda x(f)(n)fx$.

The relation of *execution* $>$ is the least preorder on $\Lambda \star \Pi$ defined by the following rules (with $\xi, \eta, \zeta \in \Lambda, \pi \in \Pi$ and $n \in \mathbb{N}$) :

1. $(\xi)\eta \star \pi > \xi \star \eta \bullet \pi$; (*push*)
2. $B \star \xi \bullet \eta \bullet \zeta \bullet \pi > \xi \star (\eta)\zeta \bullet \pi$; (*apply*)
3. $C \star \xi \bullet \eta \bullet \zeta \bullet \pi > \xi \star \zeta \bullet \eta \bullet \pi$; (*switch*)
4. $I \star \xi \bullet \pi > \xi \star \pi$; (*no operation*)
5. $K \star \xi \bullet \eta \bullet \pi > \xi \star \pi$; (*delete*)
6. $W \star \xi \bullet \eta \bullet \pi > \xi \star \eta \bullet \eta \bullet \pi$; (*copy*)
7. $cc \star \xi \bullet \pi > \xi \star k_\pi \bullet \pi$; (*save the stack*)
8. $A \star \xi \bullet \pi > \xi \star \pi_0$; (*abort*) or (*delete the stack*)
9. $\bigwedge_i \xi_i \star \underline{n} \bullet \pi > \xi_n \star \pi$; (*oracle*)

When $\xi, \eta \in \Lambda$, we set $\xi > \eta$ iff $(\forall \pi \in \Pi)(\xi \star \pi > \eta \star \pi)$.

- *Proof-like terms.*

Let PL_0 be the countable set of terms built with the constants B, C, I, K, W, cc and the application. It is the smallest possible set of *proof-like terms*.

We shall also consider the set PL of *closed terms* (i.e. with no occurrence of p, \bar{q}) which is much bigger : it contains Λ and the oracles $\bigwedge_i \xi_i$ and is therefore of cardinality 2^{\aleph_0} .

• *Execution of processes ; definition of \perp .*

For every process $\xi \star \pi$, at most one among the rules 1 to 9 applies. By iterating these rules, we obtain the *reduction* or the *execution* of the process $\xi \star \pi$. This execution stops if and only if the stack is insufficient (rules 2 to 8) or does not begin with an integer (rule 9) or else if the process has the form $p \star \omega$ or $q_i \star \omega$.

Therefore, p and q_i are *stop instructions* ; p plays a special role because :

We set $\perp = \{\xi \star \pi \in \Lambda \star \Pi ; (\exists \omega \in \Pi)(\xi \star \pi > p \star \omega)\}$.

Lemma 1. *\mathcal{B} is a coherent realizability algebra.*

\mathcal{B} is a realizability algebra :

It remains to check that $k_\pi \star \xi \bullet \omega > \xi \star \pi$, which is done by recurrence on π :

if $\pi = \pi_0$, it is rule 8 ;

if $\pi = t \bullet \rho$ we have $k_\pi \star \xi \bullet \omega = k_{t \bullet \rho} \star \xi \bullet \omega = \ell_t k_\rho \star \xi \bullet \omega > (k_\rho)(\xi)t \star \omega > k_\rho \star \xi t \bullet \omega > \xi t \star \rho$ (recurrence hypothesis) $> \xi \star t \bullet \rho$.

\mathcal{B} is coherent :

If $\theta \in \text{PL}$ then $\theta \star \pi_0 \notin \perp$; indeed, p does not appear during the execution of $\theta \star \pi_0$.

Q.E.D.

Models and functionals

A coherent realizability algebra is useful in order to give *truth values* to formulas of ZF. In fact, we use a theory called ZF_ε [6] which is a conservative extension of ZF.

This theory has an additional strong membership relation symbol ε *which is not extensional*.

For each closed formula F of ZF_ε , we define *two truth values*, denoted $\|F\|$ and $|F|$, with $\|F\| \subset \Pi$ and $|F| \subset \Lambda$, with the relation $\xi \in |F| \Leftrightarrow (\forall \pi \in \|F\|)(\xi \star \pi \in \perp)$.

The relation $\xi \in |F|$ is also written $\xi \Vdash F$ and reads “*the term ξ realizes the formula F* ”.

All the necessary definitions are given in [5, 6, 7].

The following lemma 2 is a useful property of the BBC realizability algebra \mathcal{B} .

Lemma 2. *For all formulas A, B of ZF_ε , and all terms $\xi \in \Lambda$, we have :*

$\xi \Vdash A \rightarrow B$ iff $(\forall \eta \in \Lambda)(\eta \Vdash A \Rightarrow \xi \eta \Vdash B)$.

Indeed, by the general definition of \Vdash , we have :

$$(\xi \Vdash A \rightarrow B) \Leftrightarrow (\forall \eta \Vdash A)(\forall \pi \in \|B\|)(\xi \star \eta \bullet \pi \in \perp).$$

Now, by the above definition of \perp , it is clear that $(\xi \star \eta \bullet \pi \in \perp) \Leftrightarrow (\xi \eta \star \pi \in \perp)$ from which the result follows.

Q.E.D.

Classical realizability is an extension of forcing. As in forcing, we start with an ordinary model \mathcal{M} of ZFC (or even $ZF + V = L$) which we call the *ground model*, and we build a *realizability model* \mathcal{N} which satisfies ZF_ε in the following sense :

\mathcal{M} and \mathcal{N} have the same domain, but neither the same language, nor the same truth values. The language of \mathcal{N} has the additional binary symbol ε of *strong membership*. The truth

values of \mathcal{N} are not 0, 1 as for \mathcal{M} , but are taken in $\mathcal{P}(\mathbf{\Pi})$ endowed with a suitable structure of Boolean algebra [5, 7]. We say that \mathcal{N} *satisfies a formula* F iff there is a proof-like term θ which realizes F or equivalently, if the truth value of F is the unit of the Boolean algebra $\mathcal{P}(\mathbf{\Pi})$.

A *functional* on the ground model \mathcal{M} is a formula $F(\vec{x}, y)$ of ZF with parameters in \mathcal{M} , such that $\mathcal{M} \models \forall \vec{x} \exists! y F(\vec{x}, y)$. Denoting such a functional by f , we write $y = f(\vec{x})$ for $F(\vec{x}, y)$.

Since \mathcal{M} and \mathcal{N} have the same domain, all the functionals defined on \mathcal{M} are also defined on \mathcal{N} and *they satisfy the same equations* and even the same *Horn formulas* i.e. formulas of the form $\forall \vec{x} (f_1(\vec{x}) = g_1(\vec{x}), \dots, f_n(\vec{x}) = g_n(\vec{x}) \rightarrow f(\vec{x}) = g(\vec{x}))$.

A particularly useful binary functional on \mathcal{M} (and thus also on \mathcal{N}) is the *application*, denoted by app , which is defined as follows : $\text{app}(f, x) = \{y ; (x, y) \in f\}$.

We shall often write $f[x]$ for $\text{app}(f, x)$. This allows to consider each set in \mathcal{M} (and in \mathcal{N}) as a unary functional.

Remark. We can define a set f in \mathcal{M} by giving $f[x]$ for every x , provided that there exists a set X such that $f[x] = \emptyset$ for all $x \notin X$: take $f = \bigcup_{x \in X} \{x\} \times f[x]$.

In the ground model \mathcal{M} , every function is defined in this way but in general, *this is false in \mathcal{N}* .

Quantifiers restricted to \mathbb{N}

In [7], we defined the quantifier $\forall x^{\text{int}}$, by setting :

$\|\forall x^{\text{int}} F[x]\| = \bigcup_{n \in \mathbb{N}} \|\underline{n}\} \rightarrow F[n]\| = \{\underline{n} \bullet \pi ; n \in \mathbb{N}, \pi \in \|F[n]\|\}$, so that we have :

$\xi \Vdash \forall x^{\text{int}} F[x] \Leftrightarrow \xi \underline{n} \Vdash F[n]$ for all $n \in \mathbb{N}$;

and it is shown that it is a correct definition of the restricted quantifier to \mathbb{N} .

Indeed the equivalence $\forall x^{\text{int}} F[x] \leftrightarrow \forall x (\text{int}[x] \rightarrow F[x])$ is realized by a closed λ -term independent of F , called a *storage operator*.

The formula $\text{int}[x]$ is any formula of ZF which says that x is an integer.

Theorem 3. *If we take PL as the set of proof-like terms, and if the ground model \mathcal{M} is transitive and countable, then there exists a countable realizability model \mathcal{N} which has only standard integers, i.e. which is an ω -model.*

Let \mathcal{T} be the theory formed with closed formulas, with parameters in \mathcal{M} , which are realized by a proof-like term. \mathcal{T} is ω -complete : indeed, if $\theta_n \in \text{PL}$ and $\theta_n \Vdash F[n]$ for $n \in \mathbb{N}$, let us set $c = \bigwedge_i \theta_i$. Then $c \underline{n} \Vdash F[n]$ for all $n \in \mathbb{N}$ and therefore $c \Vdash \forall n^{\text{int}} F[n]$, i.e. $\forall n^{\text{int}} F[n] \in \mathcal{T}$.

It follows that \mathcal{T} has a countable ω -model.

Q.E.D.

Proposition 4.

Let $f : \mathbb{N} \rightarrow 2$ and $\theta \in \text{PL}$, $\theta \Vdash \exists n^{\text{int}} (f(n) = 1)$. Then $\theta \star \rho \bullet \pi_0 \succ \rho \star \underline{n} \bullet \omega$ with $f(n) = 1$.

There exists $\tau \in \mathbf{\Lambda}$ such that $\tau \underline{n} \succ \rho$ if $f(n) = 1$ and $\tau \underline{n} \succ \rho_0$ if $f(n) = 0$: set $\tau = \lambda x (\bigwedge_i \xi_i) x \rho \rho_0$ with $\xi_n = \text{K}$ if $f(n) = 1$ and $\xi_n = \text{Kl}$ if $f(n) = 0$.

Then we have $\tau \Vdash \forall n^{\text{int}} (f(n) \neq 1)$ and therefore $\theta \tau \Vdash \perp$. We necessarily have :

$\theta \star \tau \bullet \pi_0 \succ \tau \star \underline{n} \bullet \pi$ for some n ; furthermore, we have $\tau \underline{n} \succ \rho$, otherwise we should have $\tau \underline{n} \succ \rho_0$, and thus $\theta \star \tau \bullet \pi_0 \notin \perp$. Therefore $f(n) = 1$.

Q.E.D.

Remark. This shows that, from any proof-like term which realizes a given Σ_1^0 arithmetical formula, we obtain a program which computes an integer satisfying this formula. Such a realizer is given by any proof of this formula by means of axioms which have themselves such realizers.

The theory of classical realizability gives realizers for the axioms of ZF. We show below that the bar recursion operator realizes the axiom of dependent choice. Finally, in section 4, we get (rather complicated) proof-like realizers for the axioms “ \mathbb{R} is well ordered” and “Continuum hypothesis”.

Execution of processes

Notation. If $\pi = t_0 \bullet \dots \bullet t_{n-1} \bullet \pi_0$, we shall write $\pi \bullet t$ for $t_0 \bullet \dots \bullet t_{n-1} \bullet t \bullet \pi_0$. Thus, we obtain $k_{\pi \bullet t}$ by replacing, in k_π , the last occurrence of A by $\ell_t A$.

Lemma 5. *If $\xi \star \pi \in \perp$, then $\xi' \star \pi' \in \perp$ and $\xi' \star \pi' \bullet t \in \perp$, where $\xi' \star \pi'$ is obtained by replacing, in $\xi \star \pi$, some occurrences of A by $(\ell_u)A = k_{u \bullet \pi_0}$ and some occurrences of the variables q_0, \dots, q_N by t_0, \dots, t_N ; t_0, \dots, t_N, t, u are arbitrary terms.*

Remark. In particular, it follows that $\xi \star \pi_0 \in \perp \Rightarrow \xi \Vdash \perp$.

Proof by recurrence on the length of the execution of $\xi \star \pi \in \perp$ by means of rules 1 to 9. We consider the last used rule. There are two non trivial cases :

- Rule 7 (execution of cc) ; we must show $cc \star \xi' \bullet \pi' \bullet t \in \perp$.

We apply the recurrence hypothesis to $\xi \star k_\pi \bullet \pi$, in which we replace :

- π_0 by $t \bullet \pi_0$ (thus π becomes $\pi \bullet t$) ;
- the last occurrence of A in $k_\pi = (\ell_{t_0}) \dots (\ell_{t_{n-1}})A$ by $(\ell_t)A$ (thus k_π becomes $k_{\pi \bullet t}$).

Then, we make the substitutions in ξ, π , which gives $\xi' \star k_{\pi' \bullet t} \bullet \pi' \bullet t$.

- Rule 8 (execution of A) ; we must show $(\ell_u)A \star \xi' \bullet \pi' \bullet t \in \perp$.

We apply the recurrence hypothesis to $\xi \star \pi_0$, which gives $\xi' \star u \bullet \pi_0 \in \perp$, thus $\xi' u \star \pi_0 \in \perp$ and therefore $A \star \xi' u \bullet \pi' \bullet t \in \perp$ (rule 8) ; finally, we obtain $(\ell_u)A \star \xi' \bullet \pi' \bullet t \in \perp$.

Q.E.D.

In each process $\xi \star \pi \in \perp$, we define an occurrence of p , which is called *efficient*, by recurrence on the length of its reduction. If $\xi = p$, it is this very occurrence. Otherwise, we consider the last rule used in the reduction, and the definition is clear ; for example, if it is rule 7, and if the efficient occurrence in $\xi \star k_\pi \bullet \pi$ is in k_π or in π , then we take the corresponding occurrence in $cc \star \xi \bullet \pi$.

Lemma 6. *If $\xi \star \pi \in \perp$, then :*

- $\xi' \star \pi' \in \perp$, where $\xi' \star \pi'$ is obtained by substituting arbitrary terms for the non efficient occurrences of p .
- $\xi' \star \pi' \notin \perp$ and indeed $\xi' \star \pi' > q_0 \star \omega$, where $\xi' \star \pi'$ is obtained by substituting q_0 for the efficient occurrence of p , and arbitrary terms for the non efficient occurrences of p .

The proof is immediate, by recurrence on the length of the reduction of $\xi \star \pi$ by means of rules 1 to 9 : consider the last used rule.

Q.E.D.

Corollary 7.

If $\xi \Vdash \top, \perp \rightarrow \perp$ and $\xi \Vdash \perp, \top \rightarrow \perp$, then $\xi \Vdash \top, \top \rightarrow \perp$, and thus :

$\lambda x(x) \Vdash \neg \forall x^{\mathbb{J}2}(x \neq 0, x \neq 1 \rightarrow \perp)$ and $W \Vdash \forall x^{\mathbb{J}2}(\forall y^{\mathbb{J}2}(y \neq 0, y \neq x \rightarrow y \not\leq x), x \neq 0 \rightarrow \perp)$.

Remark. These two formulas express respectively that the Boolean algebra $\mathbb{J}2$ is non trivial and that it is atomless.

We apply lemma 6 to $\xi \star \mathbf{p} \cdot \mathbf{p} \cdot \pi_0$. We have $\xi \star \mathbf{q}_0 \cdot \mathbf{p} \cdot \pi_0 \in \perp$ and $\xi \star \mathbf{p} \cdot \mathbf{q}_0 \cdot \pi_0 \in \perp$, which shows that the efficient occurrence of \mathbf{p} is in ξ . Therefore $\xi \star t \cdot u \cdot \pi_0 \in \perp$ for every $t, u \in \Lambda$, again by lemma 6.

The last two assertions follow from the fact that :

$\|\forall x^{\mathbb{J}2}(x \neq 0, x \neq 1 \rightarrow \perp)\| = \|\top, \perp \rightarrow \perp\| \cup \|\perp, \top \rightarrow \perp\|$ and therefore :

$|\forall x^{\mathbb{J}2}(x \neq 0, x \neq 1 \rightarrow \perp)| = |\top, \top \rightarrow \perp|$.

Q.E.D.

Theorem 8. For every sequence $\xi_i \in \Lambda$ ($i \in \mathbb{N}$), there exists $\phi \in \Lambda$ such that :

- $\phi \underline{i} > \xi_i$ for every $i \in \mathbb{N}$;
- for every $U \in \Lambda$ such that $U\phi \Vdash \perp$, there exists $k \in \mathbb{N}$ such that $U\psi \Vdash \perp$ for every $\psi \in \Lambda$ such that $\psi \underline{i} > \xi_i$ for every $i < k$.

Remark. Theorem 8 will be used in order to show properties of the bar recursion operator. In fact, the following weaker formulation is sufficient :

For every sequence $\xi_i \in \Lambda$ ($i \in \mathbb{N}$) and every $U \in \Lambda$ such that :

$$(\forall k \in \mathbb{N})(\exists \psi \in \Lambda)\{U\psi \not\Vdash \perp, (\forall i < k)(\psi \underline{i} > \xi_i)\}$$

there exists $\phi \in \Lambda$ such that $U\phi \not\Vdash \perp$ and $(\forall i \in \mathbb{N})(\phi \underline{i} > \xi_i)$.

In the particular case of forcing, this is exactly the *decreasing chain condition* : every decreasing sequence of (non false) conditions has a lower bound (which is non false).

We set $\eta_i = \lambda \mathbf{p} \lambda \mathbf{q} \vec{\xi}_i$; thus, we have $\eta_i \in \text{PL}$ and $\eta_i \mathbf{p} \vec{\mathbf{q}} > \xi_i$. Let $\eta = \bigwedge_i \eta_i$ and $\phi = \lambda x(\eta)x \mathbf{p} \vec{\mathbf{q}}$.

Thus, we have $\eta \in \text{PL}$ and $\phi \underline{i} > \xi_i$. We may assume that the constant η does not appear in U .

By the inductive definition of \bigwedge_i , it does not appear either in ξ_i .

Now, we have $U\phi \Vdash \perp \Leftrightarrow U \star \phi \cdot \pi_0 \in \perp$ (lemma 5). During the execution of the process $U \star \phi \cdot \pi_0$, the constant η arrives in head position a finite number of times, always through ϕ (since it is deleted each time it arrives in head position), therefore as follows :

$$\eta \star \underline{i} \cdot \mathbf{p} \cdot \vec{\mathbf{q}} \cdot \pi > \xi_i \star \pi.$$

Let k be an integer, greater than all the arguments of η during this execution and let $\psi \in \Lambda$ be such that $\psi \underline{i} > \xi_i$ for all $i < k$. Let us set $\tau = \lambda x \lambda \mathbf{p} \lambda \mathbf{q} \vec{\psi} x$; thus, we have $\tau \underline{i} \mathbf{p} \vec{\mathbf{q}} > \psi \underline{i} > \xi_i$ for $i < k$. In the process $U \star \phi \cdot \pi_0$, let us replace the constant η by the term τ ; we obtain $U \star \psi \cdot \pi_0$. The execution is the same, and therefore $U \star \psi \cdot \pi_0 \in \perp$ and $U\psi \Vdash \perp$.

Q.E.D.

The bar recursion operator

We define below two *proof-like terms* χ and Ψ (which are, in fact, closed λ -terms).

In these definitions, the variables i, k represent (intuitively) integers and the variable f represents a function of domain \mathbb{N} , with arbitrary values in Λ .

- We want a λ -term χ such that :

$$\chi k f z \underline{i} > f \underline{i} \text{ if } i < k ; \chi k f z \underline{i} > z \text{ if } i \geq k.$$

Therefore, we set :

$$\chi = \lambda k \lambda f \lambda z \lambda i ((i < k)(f) i) z$$

where the boolean $(i < k)$ is defined by :

$$(i < k) = ((k A) \lambda d \mathbf{0})(i A) \lambda d \mathbf{1}$$

with $\mathbf{0} = \lambda x \lambda y y$ or KI , $\mathbf{1} = \lambda x \lambda y x$ or K and $A = \lambda x \lambda y y x$ or CI .

The term $\chi \underline{k} f$ is a representation, in λ -calculus, of the finite sequence $(f \underline{0}, f \underline{1}, \dots, f \underline{k-1})$.

• We want a λ -term Ψ such that :

$$\Psi g u \underline{k} f > (u)(\chi \underline{k} f)(g) \lambda z (\Psi g u \underline{k}^+) (\chi) \underline{k} f z$$

where $k^+ = ((BW)(C)(B)BB)k$ or $\lambda f \lambda x (f)(k) f x$ is the successor of the integer k . Thus, we set :

$$\Psi = \lambda g \lambda u (Y) \lambda h \lambda k \lambda f (u)(\chi \underline{k} f)(g) \lambda z (h k^+) (\chi) \underline{k} f z.$$

where Y is the Turing fix point operator : $Y = XX$ with $X = \lambda x \lambda f (f)(x) x f$ or $(W)(B)(BW)(C)B$.

The term Ψ will be called the *bar recursion operator*.

2 Realizing countable choice

The *axiom of countable choice* is the following formula :

$$(CC) \quad \forall n \exists x F[n, x] \rightarrow \exists f \forall n^{\text{int}} F[n, f[n]]$$

where $F[n, x]$ is an arbitrary formula of ZF_ε (see [6]), with parameters and two free variables. The notation $f[n]$ stands for $\text{app}(f, n)$ (the functional app has been defined above).

Remark. This is a strong form of countable choice which shows that, in the realizability model \mathcal{N} , every countable sequence has the form $n \mapsto f[n]$ for some f . This will be used in section 4.

Theorem 9. $\lambda g \lambda u (\Psi) g u \underline{0} \underline{0} \Vdash CC$.

The axiom of countable choice is therefore realized in the model of ZF associated with the BBC realizability algebra (in fact, it is sufficient that the realizability algebra satisfies the property formulated in the remark following theorem 8).

We write the axiom of countable choice as follows :

$$(CC) \quad \forall n \neg \forall x \neg F[n, x], \forall f \neg \forall n^{\text{int}} F[n, f[n]] \rightarrow \perp$$

Let $G, U \in \mathbf{\Lambda}$ be such that $G \Vdash \forall n \neg \forall x \neg F[n, x]$ and $U \Vdash \forall f \neg \forall n^{\text{int}} F[n, f[n]]$.

We set $H = \Psi G U$ and we have to show that $H \underline{0} \underline{0} \Vdash \perp$. In fact, we shall show that $H \underline{0} \xi \Vdash \perp$ for every $\xi \in \mathbf{\Lambda}$.

Lemma 10. Let $k \in \mathbb{N}$ and $\phi \in \mathbf{\Lambda}$ be such that $(\forall i < k) \exists a_i (\phi \underline{i} \Vdash F[i, a_i])$.

If $H \underline{k} \phi \not\Vdash \perp$, then there exist a set a_k and a term $\zeta_{k, \phi} \in \mathbf{\Lambda}$ such that :

$$\zeta_{k, \phi} \Vdash F[k, a_k] \text{ and } (H \underline{k}^+) (\chi) \underline{k} \phi \zeta_{k, \phi} \not\Vdash \perp.$$

Define $\eta_{k, \phi} = \lambda z (H \underline{k}^+) (\chi) \underline{k} \phi z$, so that $H \underline{k} \phi > (U)(\chi \underline{k} \phi)(G) \eta_{k, \phi}$.

If $\eta_{k, \phi} \Vdash \forall x \neg F[k, x]$ then, by hypothesis on G , we have $G \eta_{k, \phi} \Vdash \perp$. Let us check that :

$$(\chi \underline{k} \phi)(G) \eta_{k, \phi} \Vdash \forall n^{\text{int}} F[n, f_k[n]]$$

where f_k is defined by : $f_k[i] = a_i$ if $i < k$ (i.e. $i \in k$) ; $f_k[i] = \emptyset$ if $i \notin k$.

Indeed, if we set $\phi' = (\chi \underline{k} \phi)(G) \eta_{k, \phi}$, we have :

$\phi' \underline{i} > \phi \underline{i} \Vdash F[i, a_i]$ for $i < k$ and $\phi' \underline{i} > (G) \eta_{k, \phi} \Vdash \perp$ for $i \geq k$, and therefore $\phi' \underline{i} \Vdash F[i, \emptyset]$.

By hypothesis on U , it follows that $(U)(\chi \underline{k} \phi)(G)\eta_{k,\phi} \Vdash \perp$, in other words $H\underline{k}\phi \Vdash \perp$. Thus, we have shown that, if $H\underline{k}\phi \not\Vdash \perp$, then $\eta_{k,\phi} \not\Vdash \forall x \neg F[k, x]$, which gives immediately the desired result.

Q.E.D.

Let $\phi_0 \in \Lambda$ be such that $H\underline{0}\phi_0 \not\Vdash \perp$. By means of lemma 10, we define $\phi_{k+1} \in \Lambda$ and a_k recursively on k , by setting $\phi_{k+1} = \chi \underline{k} \phi_k \zeta_{k,\phi_k}$.

By definition of χ , we have $\phi_{k+1} \underline{i} > \zeta_{k,\phi_k}$ for $i \geq k$.

Then, we show easily, by recurrence on k :

$$\phi_{k+1} \underline{i} > \phi_{i+1} \underline{i} > \zeta_{i,\phi_i} \Vdash F[i, a_i] \text{ for } i \leq k; H\underline{k}\phi_k \not\Vdash \perp.$$

Therefore, we can define:

a function f of domain \mathbb{N} such that $f[i] = a_i$ for every $i \in \mathbb{N}$;

and, by theorem 8, a term $\phi \in \Lambda$ such that $\phi \underline{k} > \zeta_{k,\phi_k}$ for all $k \in \mathbb{N}$.

Therefore, we have $\phi \underline{i} \Vdash F[i, f[i]]$ for every $i \in \mathbb{N}$, that is to say $\phi \Vdash \forall n^{\text{int}} F[n, f[n]]$.

By hypothesis on U , it follows that $U\phi \Vdash \perp$. Therefore, by theorem 8, applied to the sequence $\xi_i = \zeta_{i,\phi_i}$, there exists an integer k such that $U\psi \Vdash \perp$, for every term $\psi \in \Lambda$ such that $\psi \underline{i} > \zeta_{i,\phi_i}$ for $i < k$.

Thus, in particular, we have $(U)(\chi \underline{k} \phi_k)\xi \Vdash \perp$ for every $\xi \in \Lambda$.

Now, by definition of H , we have $H\underline{k}\phi_k > (U)(\chi \underline{k} \phi_k)\xi$ with $\xi = (G)\lambda z(H\underline{k}^+)(\chi)\underline{k}\phi_k z$, and therefore $H\underline{k}\phi_k \Vdash \perp$, that is a contradiction.

Thus, we have shown that $H\underline{0}\phi_0 \Vdash \perp$ for every $\phi_0 \in \Lambda$.

Q.E.D.

3 Realizing dependent choice

The *axiom of dependent choice* is the following formula:

$$(DC) \quad \forall x \exists y F[x, y] \rightarrow \exists f \forall n^{\text{int}} F[f[n], f[n+1]]$$

where $F[x, y]$ is an arbitrary formula of ZF_ε , with parameters and two free variables.

The notation $f[n]$ stands for $\text{app}(f, n)$ as defined above.

Theorem 11. $\lambda g \lambda u (\Psi) g u \underline{0} \underline{0} \Vdash DC$.

The axiom of dependent choice is therefore realized in the model of ZF associated with the BBC realizability algebra (or, more generally, with any realizability algebra satisfying the property formulated in the remark after theorem 8).

The proof of theorem 11 is almost the same as theorem 9.

We write the *axiom of dependent choice* as follows:

$$(DC) \quad \forall x \neg \forall y \neg F[x, y], \forall f \neg \forall n^{\text{int}} F[f[n], f[n+1]] \rightarrow \perp.$$

Let $G, U \in \Lambda$ be such that $G \Vdash \forall x \neg \forall y \neg F[x, y]$ and $U \Vdash \forall f \neg \forall n^{\text{int}} F[f[n], f[n+1]]$.

We set $H = \Psi G U$ and we have to show that $H \underline{0} \underline{0} \Vdash \perp$. In fact, we shall show that $H \underline{0} \xi \Vdash \perp$ for every $\xi \in \Lambda$.

Lemma 12.

Let a_0, \dots, a_k be a finite sequence in \mathcal{M} and $\phi \in \Lambda$ be such that $(\forall i < k)(\phi \underline{i} \Vdash F[a_i, a_{i+1}])$.

If $H\underline{k}\phi \not\Vdash \perp$, then there exist $\zeta \in \Lambda$ and a_{k+1} in \mathcal{M} such that:

$$\zeta \Vdash F[a_k, a_{k+1}] \text{ and } (H\underline{k}^+)(\chi)\underline{k}\phi\zeta \not\Vdash \perp.$$

Define $\eta_{k,\phi} = \lambda z(H\underline{k}^+)(\chi)\underline{k}\phi z$, so that $H\underline{k}\phi > (U)(\chi\underline{k}\phi)(G)\eta_{k,\phi}$.

If $\eta_{k,\phi} \Vdash \forall y \neg F[a_k, y]$ then, by hypothesis on G , we have $G\eta_{k,\phi} \Vdash \perp$. We check that :

$$(\chi\underline{k}\phi)(G)\eta_{k,\phi} \Vdash \forall n^{\text{int}} F[f_k(n), f_k[n+1]]$$

where f_k is defined by $f_k[i] = a_i$ for $i \leq k$ (i.e. $i \in k+1$) ; $f_k[i] = \emptyset$ for $i \notin k+1$.

Indeed, if we set $\phi' = (\chi\underline{k}\phi)(G)\eta_{k,\phi}$, we have :

$\phi' \underline{i} > \phi \underline{i} \Vdash F[a_i, a_{i+1}]$ for $i < k$ and $\phi' \underline{i} > (G)\eta_{k,\phi} \Vdash \perp$ for $i \geq k$.

Therefore, we have $\phi' \underline{i} \Vdash F[f_k[i], f_k[i+1]]$ for every $i \in \mathbb{N}$.

By hypothesis on U , it follows that $(U)(\chi\underline{k}\phi)(G)\eta_{k,\phi} \Vdash \perp$, that is $H\underline{k}\phi \Vdash \perp$.

Thus, we have shown that, if $H\underline{k}\phi \not\Vdash \perp$, then $\eta_{k,\phi} \not\Vdash \forall y \neg F[a_k, y]$, which gives immediately the desired result.

Q.E.D.

Let $\phi_0 \in \Lambda$ be such that $H\underline{0}\phi_0 \not\Vdash \perp$ and let $a_0 = \emptyset$. Using lemma 12, we define $\phi_{k+1} \in \Lambda$ and a_{k+1} in \mathcal{M} recursively on k , by setting $\phi_{k+1} = \chi\underline{k}\phi_k \zeta_{k,\phi_k}$, where ζ_{k,ϕ_k} is given by lemma 12, where we set $\phi = \phi_k$. By definition of χ , we have $\phi_{k+1} \underline{i} > \zeta_{k,\phi_k}$ for $i \geq k$.

Then, we show easily, by recurrence on k :

$$\phi_{k+1} \underline{i} > \phi_{i+1} \underline{i} > \zeta_{i,\phi_i} \Vdash F[a_i, a_{i+1}] \text{ for } i \leq k ; H\underline{k}\phi_k \not\Vdash \perp.$$

Therefore, we can define :

a function f of domain \mathbb{N} such that $f[i] = a_i$ for every $i \in \mathbb{N}$;

and, by means of theorem 8, a term $\phi \in \Lambda$ such that $\phi \underline{k} > \zeta_{k,\phi_k}$ for every $k \in \mathbb{N}$.

Thus, we have $\phi \underline{i} \Vdash F[f[i], f[i+1]]$ for every $i \in \mathbb{N}$, that is to say $\phi \Vdash \forall n^{\text{int}} F[f[n], f[n+1]]$.

By hypothesis on U , it follows that $U\phi \Vdash \perp$. Therefore, by theorem 8, applied to the sequence $\xi_i = \zeta_{i,\phi_i}$, there exists an integer k such that $U\psi \Vdash \perp$, for every term $\psi \in \Lambda$ such that $\psi \underline{i} > \zeta_{i,\phi_i}$ for $i < k$.

Thus, in particular, we have $(U)(\chi\underline{k}\phi_k)\xi \Vdash \perp$ for every $\xi \in \Lambda$.

But, by definition of H , we have $H\underline{k}\phi_k > (U)(\chi\underline{k}\phi_k)\xi$ with $\xi = (G)\lambda z(H\underline{k}^+)(\chi)\underline{k}\phi_k z$, and therefore $H\underline{k}\phi_k \Vdash \perp$, that is a contradiction.

Thus, we have shown that $H\underline{0}\phi_0 \Vdash \perp$ for every $\phi_0 \in \Lambda$.

Q.E.D.

4 A well ordering on \mathbb{R}

In this section, we use the notations and the results of [7] and [8].

If F is a closed formula of ZF_ε , the notation $\Vdash F$ means that there exists a proof-like term $\theta \in \text{PL}_0$ (i.e. a closed λ_c -term) such that $\theta \Vdash F$.

In section 2, we have realized the axiom of countable choice (CC). We replace $F[n, x]$ with $\text{int}(n) \rightarrow F[n, x]$ and we add a parameter ϕ ; we obtain :

$$\Vdash \forall \phi (\forall n^{\text{int}} \exists x F[n, x, \phi] \rightarrow \exists f \forall n^{\text{int}} F[n, f[n], \phi])$$

for every formula $F[n, x, \phi]$ of ZF_ε .

In particular, taking $\phi \varepsilon 2^{\mathbb{N}}$ and $F[n, x, \phi] \equiv (x = \phi(n)) \wedge (x = 0 \vee x = 1)$

(i.e. $(n, x) \varepsilon \phi \wedge (x = 0 \vee x = 1)$), we find :

$$\Vdash (\forall \phi \varepsilon 2^{\mathbb{N}}) \exists f \forall n^{\text{int}} ((f[n] = \phi(n)) \wedge (f[n] = 0 \vee f[n] = 1)).$$

For any set f in the ground model \mathcal{M} , let $g = \{x; f[x] = 1\}$.

We have trivially $\Vdash \langle n \in g \rangle = \langle f[n] = 1 \rangle$.²

It follows that : $\Vdash \forall f \exists g \forall n ((f[n] = 0 \vee f[n] = 1) \rightarrow f[n] = \langle n \in g \rangle)$.

We have shown that : $\Vdash (\forall \phi \varepsilon 2^{\mathbb{N}}) \exists g \forall n^{\text{int}} (\phi(n) = \langle n \in g \rangle)$.

Now, in [8], we have built an ultrafilter $\mathcal{D} : \mathbb{2} \rightarrow 2$ on the Boolean algebra $\mathbb{2}$, with the following property : the model \mathcal{N} , equipped with the binary relations $\mathcal{D}(\langle x \in y \rangle)$, $\mathcal{D}(\langle x = y \rangle)$, is a model of ZF, denoted $\mathcal{M}_{\mathcal{D}}$, which is an elementary extension of the ground model \mathcal{M} . Moreover, $\mathcal{M}_{\mathcal{D}}$ is isomorphic to a transitive submodel of \mathcal{N} (considered as a model of ZF), which contains every ordinal of \mathcal{N} .

$\mathcal{M}_{\mathcal{D}}$ satisfies the axiom of choice, because we suppose that $\mathcal{M} \models \text{ZFC}$.

If we suppose that $\mathcal{M} \models \text{V=L}$, then $\mathcal{M}_{\mathcal{D}}$ is isomorphic to the class $L^{\mathcal{N}}$ of constructible sets of \mathcal{N} .

For every $\phi : \mathbb{N} \rightarrow 2$, we have obviously $\mathcal{D}(\phi(n)) = \phi(n)$. It follows that :

$$\Vdash (\forall \phi \varepsilon 2^{\mathbb{N}}) \exists g \forall n^{\text{int}} (\phi(n) = \mathcal{D}\langle n \in g \rangle).$$

This shows that the subset of \mathbb{N} defined by ϕ is in the model $\mathcal{M}_{\mathcal{D}}$: indeed, it is the element g of this model.

We have just shown that \mathcal{N} and $\mathcal{M}_{\mathcal{D}}$ have the same reals.

Therefore, \mathbb{R} is well ordered in \mathcal{N} , and we have : $\Vdash (\mathbb{R} \text{ is well ordered})$.

Moreover, if the ground model \mathcal{M} satisfies V=L , we have : $\Vdash (\text{every real is constructible})$.

Therefore, the continuum hypothesis is realized.

Since the models \mathcal{N} and $\mathcal{M}_{\mathcal{D}}$ have the same reals, every formula of analysis (closed formula with quantifiers restricted to \mathbb{N} or \mathbb{R}) has the same truth value in $\mathcal{M}_{\mathcal{D}}$, \mathcal{M} or \mathcal{N} .

It follows that :

For every formula F of analysis, we have $\mathcal{M} \models F$ if and only if $\Vdash F$.

In particular, we have $\Vdash F$ or $\Vdash \neg F$.

References

- [1] S. Berardi, M. Bezem and T. Coquand. *On the computational content of the axiom of choice*. J. Symb. Logic 63, 2 (1998) p. 600-622.
- [2] U. Berger and P. Oliva. *Modified bar recursion and classical dependent choice*. Proc. Logic Colloquium 2001 - Springer (2005) p. 89-107.
- [3] E. Engeler. *Algebras and combinators*. Algebra Universalis, vol. 13, 1 (1981) p. 389-392.
- [4] T. Griffin. *A formulæ-as-type notion of control*. Conf. record 17th A.C.M. Symp. on Principles of Progr. Languages (1990).
- [5] J.-L. Krivine. *Realizability algebras : a program to well order \mathbb{R}* . Logical Methods in Computer Science vol. 7, 3:02 (2011) p. 1-47.

²The notations $\mathbb{2}$ and $\langle F \rangle$ where F is a closed formula of ZF, with parameters in the realizability model \mathcal{N} , are defined in [7, 8]. $\mathbb{2}$ is called the *characteristic Boolean algebra of \mathcal{N}* . We have $\langle F \rangle \varepsilon \mathbb{2}$.

- [6] J.-L. Krivine. *Realizability algebras II : new models of ZF + DC*.
Logical Methods in Computer Science, vol. 8, 1:10 (2012) p. 1-28.
- [7] J.-L. Krivine. *Realizability algebras III: some examples*.
<http://arxiv.org/abs/1210.5065> (2013)³. To appear in Math. Struct. Comp. Sc.
- [8] J.-L. Krivine. *On the structure of classical realizability models of ZF*.
<http://arxiv.org/abs/1408.1868> (2014). To appear in Proceedings Types '2014.
- [9] C. Spector. *Provably recursive functionals of analysis : a consistency proof of analysis by an extension of principles in current intuitionistic mathematics*.
Recursive function theory : Proc. Symp. in pure math. vol. 5, Amer. Math. Soc. Providence, Rhode Island, 1962, p. 1-27.
- [10] T. Streicher. *A classical realizability model arising from a stable model of untyped λ -calculus*. <http://arxiv.org/abs/1407.1547> (2013).

³Articles [5, 6, 7, 8] are available at www.pps.univ-paris-diderot.fr/~krivine/